

References:

1. Ponikowski P, et al. ESC Heart Fail. 2014;1(1):4-25.
2. Krumholz HM, et al. Circ Cardiovasc Qual Outcomes. 2009;2:407-413.
3. Cowie MR, et al. Improving care for patients are acute heart failure [Internet; cited 2013]. Available from: <http://www.oxfordhealthpolicyforum.org/reports/acute-heart-failure/improving-care-for-patients-with-acute-heart-failure>.
4. Butler J, et al. Congest Heart Fail. 2012;18(5 Suppl 1):S1-S3.
5. Chioncel O, et al. Eur J Heart Fail. 2017;19:1242-1254;
6. Granger BB, et al., J Card Fail. 2022;28:1355-1361.
7. Butler J, et al. Eur J Heart Fail. 2021;23:1334-1342.
8. Greene SJ, et al., J Am Coll Cardiol. 2018;72:351-366.
9. Joseph S, et al. Eur Heart J. 2021;43:908-915.
10. Damasceno A, et al., Arch Intern Med. 2012;172:1386-1394.
11. Čerlinskaitė K, et al. ESC Heart Fail. 2021;8:2473-2484.
12. McDonagh TA, et al. Eur Heart J. 2021 Sep 21;42(36):3599-3726.
13. Komajda M, et al. Eur J Heart Fail. 2016;18(5):514-22.
14. Mebazaa A, et al. The Lancet. 2022; S0140-6736(22)02076-1.

All trademarks mentioned
enjoy legal protection.

© 2022 Roche

Published by
Roche Diagnostics Scandinavia AB
Box 1228
171 23 Solna
MC-SE-02154

STR♥NG-HF


CONTEMPORARY POST-DISCHARGE MANAGEMENT IN HEART-FAILURE

Primary outcomes


This material is intended for use
by healthcare professionals.

Heart failure patients currently have high rates of readmission and mortality after discharge

The burden of heart failure




What are the challenges of post-discharge management?



STRONG-HF primary outcomes show significant reduction of all cause death or acute heart failure readmissions

Study Design



Results

